



The Mantoux tuberculin skin test is the standard method of identifying persons infected with Mycobacterium Tuberculosis (TB). Tuberculin skin testing is both safe and reliable.

It is recommended that those who work in a health care setting and/or have an increased exposure risk have a two-step testing completed. A TB risk assessment questionnaire will be used yearly to identify any persons who remain at an increased risk of acquiring TB or those who exhibit symptoms of illness.

- | | | |
|---|-----|----|
| 1. Have you ever had the BCG vaccine? | Yes | No |
| 2. Are you pregnant? | Yes | No |
| 3. Have you ever had a positive TB test? | Yes | No |
| 4. Have you ever had a severe allergic reaction to PPD? | Yes | No |

By signing below I give consent to have the Mantoux skin test to screen for TB using the purified protein derivative (PPD). 0.1ml of PPD is injected intradermally in the forearm producing a pale elevation of skin (wheal) of 6-10mm.

Name (print) _____ Signature _____

Step 1:

Date Given: _____ Signatures/Title: _____ Site: _____

NDC: _____ Vaccine Lot: _____ Exp. _____

*Test must be read by a nurse or provider in 48-72 hours

Date Read: _____ Result: _____ Signature/Title: _____

Step 2:

Date Given: _____ Signatures/Title: _____ Site: _____

NDC: _____ Vaccine Lot: _____ Exp. _____

*Test must be read by an RN or Medical Provider in 48-72 hours

Date Read: _____ Result: _____ Signature/Title: _____